arXiv:1206.3826v1 [math.NT] 18 Jun 2012

ON THE INTEGRAL OF THE PRODUCT OF FOUR AND MORE BERNOULLI POLYNOMIALS

DAEYEOUL KIM, SU HU, AND MIN-SOO KIM

Abstract. L.J. Mordell in [*Integral formulae of arithmetical character*, J. London Math. Soc. **33** (1958) 371–375] provided the formula for the integral of the product of two Bernoulli polynomials, he also remarked: "The integrals containing the product of more than two Bernoulli polynomials do not appear to lead to simple results." Let r be any positive integer. In this paper, we provide the explicit formulas for the integral of the product of r Bernoulli polynomials. Many authors' results in this direction, including Nörlund, Mordell, Carlitz, Agoh and Dilcher are special cases of the formulas given in this paper.

1. Introduction

Nearly 90 years ago, Nörlund gave the following formula for the integral of the product of two Bernoulli polynomials in his book "Vorlesungen uber Differenzenrechnung":

$$\int_0^1 B_k(z)B_l(z)dz = (-1)^{k-1}\frac{k!l!}{(k+l)!}B_{k+l},\ k+l\geq 2 \tag{1.1}$$

(see [8, p. 31]).

Nikolas and Mordell provided two different proofs of (1.1) in [8] and [7] respectively. Mordell remarked: " The integrals containing the product of more than two Bernoulli polynomials do not appear to lead to simple results." (See [7, p. 375]). But Carlitz [3] obtained an explicit formula for the integral of the product of three and four Bernoulli polynomials which extends (1.1). Later, Wilson [10] generalized Carlitz's result on the integral of the product of three Bernoulli polynomials by

---

2010 *Mathematics Subject Classification.* 11B68.

Key words and phrases. Bernoulli polynomials; Bernoulli numbers; Integrals; Recurrence relations.

evaluating the following integral

$$\int_0^1 \overline{B}_k(az)\overline{B}_l(bz)\overline{B}_m(cz)dz, \tag{1.2}$$

where $\overline{B}_k(x)$ is the periodic extension of $B_k(x)$ on $[0,1)$ and a,b,c are pairwise coprime integers. Carlitz's result became special case when $a=b=c=1$. Similar integral evaluations have also been used by Espinosa and Moll [5] during their study on Tornheim's double sums.

Recently, Wilson's result has been generalized by Agoh and Dilcher [2]. In fact, they proved the following result:

Proposition 1.1 ([2, Proposition 3])**.** *For* $k,l,m\geq 0$

$$\frac{I_{k,l,m}}{k!l!m!}=\sum_{a=0}^{k+l}(-1)^a\sum_{i=0}^{a}\binom{a}{i}\frac{C_{k-a+i,l-i,m+a+1}}{(k-a+i)!(l-i)!(m+a+1)!},$$

where

$$\begin{aligned}I_{k,l,m}&=\int_0^x B_k(z)B_l(z)B_m(z)dz,\\ C_{k,l,m}&=B_k(x)B_l(x)B_m(x)-B_kB_lB_m.\end{aligned}$$

Let r be any positive integer. In this paper, we provide the explicit formulas for the integral of the product of r Bernoulli polynomials which lead all the previous mentioned results for the integral of the products of Bernoulli polynomials become special cases (see Remark 1.5 below).

Before statement our main results, we introduce the definition and some basic properties on multinomial coefficients. Since the Binomials coefficients have been used by Agoh and Dilcher in [2] to get their formulas of the product of three Bernoulli polynomials, in order to get more general ones, we need some information on multinomial coefficients. They are defined to be the number:

$$\binom{\mu}{k_1,\ldots,k_r}=\frac{\mu!}{k_1!\cdots k_r!}, \tag{1.3}$$

where $k_1+\cdots+k_r=\mu$ and $k_1,\ldots,k_r\geq 0$. When $r=2$, we obtain the binomial coefficients

$$\binom{\mu}{k_1,k_2}=\binom{\mu}{k_1,\mu-k_1}=\binom{\mu}{k_1}=\binom{\mu}{k_2}. \tag{1.4}$$

The multinomial coefficient can be extended as follows

$$\binom{\mu}{k_1,\dots,k_r} = \begin{cases} 0 & \text{if } \min\limits_{1\le i\le r}\{k_i\} < 0 \text{ and } \max\limits_{1\le i\le r}\{k_i\} > \mu \\ \frac{\mu!}{k_1!\cdots k_r!} & \text{otherwise.} \end{cases} \tag{1.5}$$

Lemma 1.2. *Multinomial coefficients have the recurrence relation*

$$\binom{\mu}{k_1,\dots,k_r} = \binom{\mu-1}{k_1-1,\dots,k_r} + \cdots + \binom{\mu-1}{k_1,\dots,k_r-1}$$

and symmetry

$$\binom{\mu}{k_1,\dots,k_r} = \binom{\mu}{k_{\sigma(1)},\dots,k_{\sigma(r)}},$$

where σ is a permutation of $(1,\dots,r)$.

The following two are the main results of this paper.

Theorem 1.3. *Let*

$$\begin{aligned} I_{k_1,\dots,k_r} &= \int_0^x B_{k_1}(z)\cdots B_{k_r}(z)dz, \\ C_{k_1,\dots,k_r} &= B_{k_1}(x)\cdots B_{k_r}(x) - B_{k_1}\cdots B_{k_r}, \\ \widetilde{I}_{k_1,\dots,k_r} &= \frac{1}{k_1!\cdots k_r!} I_{k_1,\dots,k_r}, \\ \widetilde{C}_{k_1,\dots,k_r} &= \frac{1}{k_1!\cdots k_r!} C_{k_1,\dots,k_r}. \end{aligned} \tag{1.6}$$

For any $\mu \ge 1$ and $k_1,\dots,k_r \ge 0$, we have

$$\begin{aligned} \widetilde{I}_{k_1,\dots,k_r} = \sum_{a=0}^{\mu-1}(-1)^a & \sum_{\substack{i_1+\cdots+i_{r-1}=a \\ 0\le i_1,\dots,i_{r-1}\le a}} \binom{a}{i_1,\dots,i_{r-1}} \\ & \times \widetilde{C}_{k_1-i_1,\dots,k_{r-1}-i_{r-1},k_r+a+1} \\ & + (-1)^\mu \sum_{\substack{i_1+\cdots+i_{r-1}=\mu \\ 0\le i_1,\dots,i_{r-1}\le \mu}} \binom{\mu}{i_1,\dots,i_{r-1}} \\ & \times \widetilde{I}_{k_1-i_1,\dots,k_{r-1}-i_{r-1},k_r+\mu}. \end{aligned} \tag{1.7}$$

Letting

$$\widetilde{I}_{k_1,\dots,k_r} = 0 \quad \text{when } \min_{1\le i\le r}\{k_i\} < 0. \tag{1.8}$$

and

$$\mu = k_1 + \cdots + k_{r-1} + 1$$

in Theorem 1.3, we obtain the following result.

Theorem 1.4. *For any* $k_1, \ldots, k_r \geq 0$, *we have*

$$\widetilde{I}_{k_1,\ldots,k_r} = \sum_{a=0}^{k_1+\cdots+k_{r-1}} (-1)^a \sum_{\substack{i_1+\cdots+i_{r-1}=a \\ 0\leq i_1,\ldots,i_{r-1}\leq a}} \binom{a}{i_1,\ldots,i_{r-1}} \times \widetilde{C}_{k_1-i_1,\ldots,k_{r-1}-i_{r-1},k_r+a+1}.$$

Remark 1.5. In Theorem 1.4, when $r = 2$ and $x = 1$, we obtain the results of Nörlund [9] and Mordell [7], when $r = 3$ and $x = 1$, we obtain the result of Carlitz [3], when $r = 3$, we obtain the result of Agoh and Dilcher [1].

Using Theorem 1.3, we immediately get some concrete examples for the integral of the product of four Bernoulli polynomials.

Example 1.6. (1) $I_{1,1,1,1} = \frac{3}{2}B_1^2B_2 + \frac{1}{4}B_4 - \frac{1}{4}B_2$.

(2) $I_{1,1,1,3} = \frac{3}{4}B_1^2B_4 + \frac{1}{20}B_6 - \frac{1}{8}B_4$.

(3) $I_{1,1,1,5} = \frac{1}{2}B_1^2B_6 + \frac{1}{56}B_8 - \frac{1}{12}B_6$.

(4) $I_{1,2,2,1} = \frac{1}{2}B_2^3 + \frac{1}{2}B_2B_4 + B_1^2B_4 + \frac{1}{6}B_6 - \frac{1}{6}B_4$.

(5) $I_{2,3,3,4} = -\frac{9}{70}B_2^2B_8 - \frac{1}{35}B_2B_{10} - \frac{3}{70}B_1^2B_{10} - \frac{1}{495}B_{12} + \frac{1}{140}B_{10}$.

From Theorem 1.4 and the definition of the integral $\widetilde{I}_{k_1,\ldots,k_r}$, we have the following result.

Corollary 1.7. *Let* $T_{k_1,\ldots,k_r}(x)$ *be the right-hand side of Theorem 1.4, and let* $\sigma \in S_r$, *where* S_r *is the symmetric group of degree* r. *Then for all* $k_1, \ldots, k_r \geq 0$,

$$T_{k_1,\ldots,k_r}(x) = T_{\sigma(k_1),\ldots,\sigma(k_r)}(x).$$

The following parts of this paper are organized as follows.

In section 2, we recall the definitions and some properties for the Bernoulli polynomials, in particular the well-known fact for the integral of one Bernoulli polynomial. They will be used in the proof of the main result of this paper. In sections 3 and 4, we prove Theorems 1.3 and 1.4 respectively. In the last section, by using our formulas, we derive several examples including some formulas by previous authors.

2. BERNOULLI POLYNOMIALS

The Bernoulli polynomials $B_k(x)$ are defined by the exponential generating function

$$\frac{te^{xt}}{e^t-1}=\sum_{k=0}^{\infty}B_k(x)\frac{t^k}{k!}. \tag{2.1}$$

They satisfy the following well-known identities:

$$B_k(x+1)-B_k(x)=kx^{k-1},\ k\geq 1, \tag{2.2}$$

$$\frac{\mathrm{d}}{\mathrm{d}x}B_k(x)=kB_{k-1}(x),\ k\geq 1, \tag{2.3}$$

$$\int_x^{x+1}B_k(z)dz=x^k,\ k\geq 0, \tag{2.4}$$

$$\int_x^y B_k(z)dz=\frac{B_{k+1}(y)-B_{k+1}(x)}{k+1},\ k\geq 0, \tag{2.5}$$

$$B_k(1-x)=(-1)^kB_k(x),\ k\geq 0. \tag{2.6}$$

Let $B_k=B_k(0)$ be the Bernoulli numbers. From the above definition of $B_k(x)$, we conclude that all the B_k are rational numbers. It is well-known that $B_{2k+1}=0$ for $k\geq 1$, and $B_k(x)$ are alternatively positive and negative for even k. Letting $x=0$ in (2.6), we get the following special value of the Bernoulli polynomials:

$$B_k(1)=\begin{cases}B_k & \text{for } k\neq 1\\ -B_1 & \text{for } k=1\end{cases} \tag{2.7}$$

(see [2]). From (2.4) and (2.5), we have

$$\begin{aligned}\sum_{i=0}^{n}i^k&=\sum_{i=0}^{n}\int_i^{i+1}B_k(z)dz\\&=\int_0^{n+1}B_k(z)dz\\&=\frac{B_{k+1}(n+1)-B_{k+1}}{k+1},\ k\geq 1.\end{aligned} \tag{2.8}$$

This is the formula for the integral of one Bernoulli polynomial.

3. Proof of Theorem 1.3

Using integration by parts, along with (1.6) and (2.3), we obtain

$$\begin{aligned} I_{k_1,\ldots,k_r} &= \frac{1}{k_r+1}\int_0^x B_{k_1}(z)\cdots B_{k_{r-1}}(z)\frac{\mathrm{d}}{\mathrm{d}z}B_{k_r+1}(z)dz \\ &= \frac{1}{k_r+1}\left[B_{k_1}(z)\cdots B_{k_{r-1}}(z)B_{k_r+1}(z)\right]_0^x \\ &\quad - \frac{1}{k_r+1}\int_0^x \frac{\mathrm{d}}{\mathrm{d}z}(B_{k_1}(z)\cdots B_{k_{r-1}}(z))B_{k_r+1}(z)dz \\ &= \frac{1}{k_r+1}C_{k_1,\ldots,k_{r-1},k_r+1} \\ &\quad - \frac{1}{k_r+1}\int_0^x \frac{\mathrm{d}}{\mathrm{d}z}(B_{k_1}(z)\cdots B_{k_{r-1}}(z))B_{k_r+1}(z)dz. \end{aligned} \tag{3.1}$$

From (2.3) we have

$$\begin{aligned} &\frac{\mathrm{d}}{\mathrm{d}z}(B_{k_1}(z)\cdots B_{k_{r-1}}(z)) \\ &\quad = k_1B_{k_1-1}(z)\cdots B_{k_{r-1}}(z) \\ &\qquad + \cdots + k_{r-1}B_{k_1}(z)\cdots B_{k_{r-1}-1}(z). \end{aligned} \tag{3.2}$$

From above and (1.6) we have

$$\begin{aligned} &\frac{1}{k_r+1}\int_0^x \frac{\mathrm{d}}{\mathrm{d}z}(B_{k_1}(z)\cdots B_{k_{r-1}}(z))B_{k_r+1}(z)dz \\ &= \frac{k_1}{k_r+1}\int_0^x B_{k_1-1}(z)\cdots B_{k_{r-1}}(z)B_{k_r+1}(z)dz \\ &\quad + \cdots + \frac{k_{r-1}}{k_r+1}\int_0^x B_{k_1}(z)\cdots B_{k_{r-1}-1}(z)B_{k_r+1}(z)dz \\ &= \frac{k_1}{k_r+1}I_{k_1-1,\ldots,k_{r-1},k_r+1} + \cdots + \frac{k_{r-1}}{k_r+1}I_{k_1,\ldots,k_{r-1}-1,k_r+1}, \end{aligned} \tag{3.3}$$

and substituting (3.3) to (3.1) we get

$$\begin{aligned} I_{k_1,\ldots,k_r} &= \frac{1}{k_r+1}C_{k_1,\ldots,k_{r-1},k_r+1} \\ &\quad - \frac{1}{k_r+1}(k_1I_{k_1-1,\ldots,k_{r-1},k_r+1} + \cdots + k_{r-1}I_{k_1,\ldots,k_{r-1}-1,k_r+1}). \end{aligned} \tag{3.4}$$

From (1.6) and (3.4), we have

$$\widetilde{I}_{k_1,\ldots,k_r} = \widetilde{C}_{k_1,\ldots,k_{r-1},k_r+1} - (\widetilde{I}_{k_1-1,\ldots,k_{r-1},k_r+1} + \cdots + \widetilde{I}_{k_1,\ldots,k_{r-1}-1,k_r+1}). \tag{3.5}$$

Now we prove the case when $\mu \geq 1$ in Theorem 1.3 by induction. For $\mu = 1$ it just reduces to (3.5). Now we assume that Theorem 1.3 holds for some $\mu \geq 1$. Write

$$S = \sum_{\substack{i_1+\cdots+i_{r-1}=\mu \\ 0\leq i_1,\ldots,i_{r-1}\leq\mu}} \binom{\mu}{i_1,\ldots,i_{r-1}} \widetilde{I}_{k_1-i_1,\ldots,k_{r-1}-i_{r-1},k_r+\mu}. \tag{3.6}$$

Eq. (1.7) becomes

$$\begin{aligned} \widetilde{I}_{k_1,\ldots,k_r} = & \sum_{a=0}^{\mu-1}(-1)^a \sum_{\substack{i_1+\cdots+i_{r-1}=a \\ 0\leq i_1,\ldots,i_{r-1}\leq a}} \binom{a}{i_1,\ldots,i_{r-1}} \\ & \times \widetilde{C}_{k_1-i_1,\ldots,k_{r-1}-i_{r-1},k_r+a+1} \\ & + (-1)^\mu S. \end{aligned} \tag{3.7}$$

From (3.5) we have

$$\begin{aligned} S = & \sum_{\substack{i_1+\cdots+i_{r-1}=\mu \\ 0\leq i_1,\ldots,i_{r-1}\leq\mu}} \binom{\mu}{i_1,\ldots,i_{r-1}} \widetilde{C}_{k_1-i_1,\ldots,k_{r-1}-i_{r-1},k_r+\mu+1} \\ & - \sum_{\substack{i_1+\cdots+i_{r-1}=\mu \\ 0\leq i_1,\ldots,i_{r-1}\leq\mu}} \binom{\mu}{i_1,\ldots,i_{r-1}} \widetilde{I}_{k_1-(i_1+1),\ldots,k_{r-1}-i_{r-1},k_r+\mu+1} \\ & - \cdots \\ & - \sum_{\substack{i_1+\cdots+i_{r-1}=\mu \\ 0\leq i_1,\ldots,i_{r-1}\leq\mu}} \binom{\mu}{i_1,\ldots,i_{r-1}} \widetilde{I}_{k_1-i_1,\ldots,k_{r-1}-(i_{r-1}+1),k_r+\mu+1}. \end{aligned} \tag{3.8}$$

Let $i'_j = i_j + 1$ in (3.8) for $j = 1, \ldots, r-1$. From (1.5), we obtain

$$\begin{aligned} & \binom{\mu}{i'_1-1,\ldots,i_{r-1}} = 0 \quad \text{if } i'_1 = 0, \\ & \ldots, \\ & \binom{\mu}{i_1,\ldots,i'_{r-1}-1} = 0 \quad \text{if } i'_{r-1} = 0, \end{aligned} \tag{3.9}$$

and

$$\begin{aligned}
&\binom{\mu}{i_1'-1,\dots,i_{r-1}} = 0 \quad \text{if } i_2 = \mu+1,\dots,i_{r-1} = \mu+1,\\
&\cdots,\\
&\binom{\mu}{i_1,\dots,i_{r-1}'-1} = 0 \quad \text{if } i_1 = \mu+1,\dots,i_{r-2} = \mu+1,
\end{aligned}\tag{3.10}$$

then we have

$$\begin{aligned}
&\sum_{\substack{i_1+\cdots+i_{r-1}=\mu\\ 0\le i_1,\dots,i_{r-1}\le\mu}} \binom{\mu}{i_1,\dots,i_{r-1}} \widetilde{I}_{k_1-(i_1+1),\dots,k_{r-1}-i_{r-1},k_r+\mu+1}\\
&= \sum_{\substack{i_1'+\cdots+i_{r-1}=\mu+1\\ 0\le i_1',\dots,i_{r-1}\le\mu+1}} \binom{\mu}{i_1'-1,\dots,i_{r-1}} \widetilde{I}_{k_1-i_1',\dots,k_{r-1}-i_{r-1},k_r+\mu+1},\\
&\cdots,\\
&\sum_{\substack{i_1+\cdots+i_{r-1}=\mu\\ 0\le i_1,\dots,i_{r-1}\le\mu}} \binom{\mu}{i_1,\dots,i_{r-1}} \widetilde{I}_{k_1-i_1,\dots,k_{r-1}-(i_{r-1}+1),k_r+\mu+1}\\
&= \sum_{\substack{i_1+\cdots+i_{r-1}'=\mu+1\\ 0\le i_1,\dots,i_{r-1}'\le\mu+1}} \binom{\mu}{i_1,\dots,i_{r-1}'-1} \widetilde{I}_{k_1-i_1,\dots,k_{r-1}-i_{r-1}',k_r+\mu+1}.
\end{aligned}\tag{3.11}$$

From (3.8) and (3.11) we get

$$\begin{aligned}
S = &\sum_{\substack{i_1+\cdots+i_{r-1}=\mu\\ 0\le i_1,\dots,i_{r-1}\le\mu}} \binom{\mu}{i_1,\dots,i_{r-1}} \widetilde{C}_{k_1-i_1,\dots,k_{r-1}-i_{r-1},k_r+\mu+1}\\
&- \sum_{\substack{i_1'+\cdots+i_{r-1}=\mu+1\\ 0\le i_1',\dots,i_{r-1}\le\mu+1}} \binom{\mu}{i_1'-1,\dots,i_{r-1}} \widetilde{I}_{k_1-i_1',\dots,k_{r-1}-i_{r-1},k_r+\mu+1}\\
&- \cdots\\
&- \sum_{\substack{i_1+\cdots+i_{r-1}'=\mu+1\\ 0\le i_1,\dots,i_{r-1}'\le\mu+1}} \binom{\mu}{i_1,\dots,i_{r-1}'-1} \widetilde{I}_{k_1-i_1,\dots,k_{r-1}-i_{r-1}',k_r+\mu+1}.
\end{aligned}\tag{3.12}$$

Let $i_j = i'_j$ for $j = 1, \ldots, r-1$ in (3.12). From Lemma 1.2, (3.12) becomes

(3.13)

$$\begin{aligned} S = &\sum_{\substack{i_1+\cdots+i_{r-1}=\mu \\ 0\le i_1,\ldots,i_{r-1}\le\mu}} \binom{\mu}{i_1,\ldots,i_{r-1}} \widetilde{C}_{k_1-i_1,\ldots,k_{r-1}-i_{r-1},k_r+\mu+1} \\ &- \sum_{\substack{i_1+\cdots+i_{r-1}=\mu+1 \\ 0\le i_1,\ldots,i_{r-1}\le\mu+1}} \left[\binom{\mu}{i_1-1,\ldots,i_{r-1}} + \cdots + \binom{\mu}{i_1,\ldots,i_{r-1}-1}\right] \\ &\times \widetilde{I}_{k_1-i_1,\ldots,k_{r-1}-i_{r-1},k_r+\mu+1} \\ = &\sum_{\substack{i_1+\cdots+i_{r-1}=\mu \\ 0\le i_1,\ldots,i_{r-1}\le\mu}} \binom{\mu}{i_1,\ldots,i_{r-1}} \widetilde{C}_{k_1-i_1,\ldots,k_{r-1}-i_{r-1},k_r+\mu+1} \\ &- \sum_{\substack{i_1+\cdots+i_{r-1}=\mu+1 \\ 0\le i_1,\ldots,i_{r-1}\le\mu+1}} \binom{\mu+1}{i_1,\ldots,i_{r-1}} \widetilde{I}_{k_1-i_1,\ldots,k_{r-1}-i_{r-1},k_r+\mu+1}. \end{aligned}$$

Substituting (3.13) to (3.7), we get Theorem 1.3 when $\mu+1$ which completes our proof.

4. PROOF OF THEOREM 1.4

Letting $\widetilde{I}_{k_1-i_1,\ldots,k_{r-1}-i_{r-1},k_r+\mu} \neq 0$ in Theorem 1.3, from (1.8), we have

$$k_1 - i_1 \ge 0, \ldots, k_{r-1} - i_{r-1} \ge 0. \tag{4.1}$$

Thus

$$\begin{aligned} &(k_1 - i_1) + \cdots + (k_{r-1} - i_{r-1}) \\ &= k_1 + \cdots + k_{r-1} - (i_1 + \cdots + i_{r-1}) \\ &= k_1 + \cdots + k_{r-1} - \mu \\ &\ge 0. \end{aligned} \tag{4.2}$$

Let $\mu = k_1 + \cdots + k_{r-1} + 1$ in (4.2). We have $k_1 + \cdots + k_{r-1} - \mu < 0$. This contradicts with (4.1). Therefore when $\mu = k_1 + \cdots + k_{r-1} + 1$, one of

$$k_1 - i_1, \ldots, k_{r-1} - i_{r-1}$$

will be always negative. Thus from (1.8), any terms

$$\widetilde{I}_{k_1-i_1,\ldots,k_{r-1}-i_{r-1},k_1+\cdots+k_{r-1}+k_r+1}$$

vanishes. The result then follows from Theorem 1.3.

5. Some further consequences

Letting $r = 2$ in Theorem 1.4, we have

$$\widetilde{I}_{k_1,k_2} = \sum_{a=0}^{k_1} (-1)^a \widetilde{C}_{k_1-a,k_2+a+1}.$$

This is equivalent to the following proposition:

Proposition 5.1 ([1, Proposition 1])**.**

$$\int_0^x B_{k_1}(z)B_{k_2}(z)dz = \frac{k_1!k_2!}{(k_1+k_2+1)!} \sum_{a=0}^{k_1} (-1)^a \binom{k_1+k_2+1}{k_1-a} \times (B_{k_1-a}(x)B_{k_2+a+1}(x) - B_{k_1-a}B_{k_2+a+1}).$$

Letting $r = 3$ in Theorem 1.4, we have:

Proposition 5.2 ([1, Proposition 3])**.**

$$\begin{aligned}\widetilde{I}_{k_1,k_2,k_3} &= \sum_{a=0}^{k_1+k_2} (-1)^a \sum_{\substack{i_1+i_2=a \\ 0\le i_1,i_2\le a}} \binom{a}{i_1, i_2} \widetilde{C}_{k_1-i_1,k_2-i_2,k_3+a+1} \\ &= \sum_{a=0}^{k_1+k_2} (-1)^a \sum_{i=0}^{a} \binom{a}{i} \frac{C_{k_1-a+i,k_2-i,k_3+a+1}}{(k_1-a+i)!(k_2-i)!(k_3+a+1)!}.\end{aligned}$$

Letting $r = 4$ in Theorem 1.4, we have:

Proposition 5.3.

$$\widetilde{I}_{k_1,k_2,k_3,k_4} = \sum_{a=0}^{k_1+k_2+k_3} (-1)^a \sum_{\substack{i_1+i_2+i_3=a \\ 0\le i_1,i_2,i_3\le a}} \binom{a}{i_1, i_2, i_3} \widetilde{C}_{k_1-i_1,k_2-i_2,k_3-i_3,k_4+a+1}.$$

Let

$$I_{k_1,k_2,k_3,k_4}(1) = \int_0^1 B_{k_1}(z)B_{k_2}(z)B_{k_3}(z)B_{k_4}(z)dz.$$

Write

$$\widetilde{B}_k = \frac{1}{k!}B_k.$$

Using (1.6), (2.6) and Proposition 5.3, we reobtain the following known evaluation of the integral from 0 to 1:

$$\begin{aligned}\frac{I_{k_1,k_2,k_3,k_4}(1)}{k_1!k_2!k_3!k_4!} = \sum_{a=0}^{k_1+k_2+k_3} &(-1)^a \sum_{\substack{i_1+i_2+i_3=a\\0\le i_1,i_2,i_3\le a}} \binom{a}{i_1,i_2,i_3} \\ &\times ((-1)^{k_1+k_2+k_3+k_4+1}-1) \\ &\times \widetilde{B}_{k_1-i_1}\widetilde{B}_{k_2-i_2}\widetilde{B}_{k_3-i_3}\widetilde{B}_{k_4+a+1},\end{aligned}$$

so

$$\begin{aligned}\frac{I_{k_1,k_2,k_3,k_4}(1)}{k_1!k_2!k_3!k_4!} = 2 &\sum_{a=0}^{k_1+k_2+k_3} (-1)^{a+1}\widetilde{B}_{k_4+a+1} \\ &\times \sum_{\substack{i_1+i_2+i_3=a\\0\le i_1,i_2,i_3\le a}} \binom{a}{i_1,i_2,i_3}\widetilde{B}_{k_1-i_1}\widetilde{B}_{k_2-i_2}\widetilde{B}_{k_3-i_3}\end{aligned} \tag{5.1}$$

if $k_1+k_2+k_3+k_4 \equiv 0 \pmod 2$, and

$$\frac{I_{k_1,k_2,k_3,k_4}(1)}{k_1!k_2!k_3!k_4!} = 0 \tag{5.2}$$

if $k_1+k_2+k_3+k_4 \not\equiv 0 \pmod 2$. Therefore, without loss of generality we assume that

$$k_1+k_2+k_3+k_4 \equiv 0 \pmod 2 \quad \text{and} \quad k_4+a+1 \equiv 0 \pmod 2 \tag{5.3}$$

since $B_{k_4+a+1} = 0$ if $k_4+a+1 \not\equiv 0 \pmod 2$. By (5.3), we see that

$$\begin{aligned}k_1+k_2+k_3+k_4+1 &= (k_1-i_1)+(k_2-i_2)+(k_3-i_3)+(k_4+a+1) \\ &\equiv 1 \pmod 2.\end{aligned} \tag{5.4}$$

Thus from (5.1) and (5.4) we get the following table:

	k_1-i_1	k_2-i_2	k_3-i_3	k_4+a+1
(A)	odd	even	even	even
(B)	even	odd	even	even
(C)	even	even	odd	even
(D)	odd	odd	odd	even

From now on, we will use the fact that $B_{2i+1} = 0$ for $i \ge 1$. Thus, if $k_j - i_j$ are odd for $j = 1,2,3$, then $k_j - i_j$ must be 1, so we only need to consider the following four cases:

Case (A). We put $i_1 = k_1 - 1$. Then by (5.3) and $a = i_1 + i_2 + i_3$, we have

$$k_2 - i_2 \equiv k_3 - i_3 \pmod 2.$$

Case (B). We put $i_2 = k_2 - 1$. Similarly, we have

$$k_1 - i_1 \equiv k_3 - i_3 \pmod 2.$$

Case (C). We put $i_3 = k_3 - 1$. Similarly, we have

$$k_1 - i_1 \equiv k_2 - i_2 \pmod 2.$$

Case (D). We have

$$i_1 = k_1 - 1, i_2 = k_2 - 1, i_3 = k_3 - 1$$

since $B_{2i+1} = 0$ for $i \geq 1$.

Using (5.1), (5.2), (A), (B), (C) and (D), we obtain the following theorem.

Theorem 5.4. (1) *If* $k_1 + k_2 + k_3 + k_4 \equiv 0 \pmod 2$, *then*

$$\begin{aligned}\frac{I_{k_1,k_2,k_3,k_4}(1)}{k_1!k_2!k_3!k_4!} = &\sum_{a=0}^{k_1+k_2+k_3} (-1)^a \widetilde{B}_{k_4+a+1} \\ &\times \left\{ \binom{a}{k_1 - 1} \sum_{i=0}^{a-k_1+1} \binom{a - k_1 + 1}{i} \widetilde{B}_{k_3-i} \widetilde{B}_{k_1+k_2+i-a-1} \right. \\ &+ \binom{a}{k_2 - 1} \sum_{i=0}^{a-k_2+1} \binom{a - k_2 + 1}{i} \widetilde{B}_{k_1-i} \widetilde{B}_{k_2+k_3+i-a-1} \\ &\left. + \binom{a}{k_3 - 1} \sum_{i=0}^{a-k_3+1} \binom{a - k_3 + 1}{i} \widetilde{B}_{k_1-i} \widetilde{B}_{k_2+k_3+i-a-1} \right\} \\ &+ \frac{1}{2}(-1)^{k_1+k_2+k_3} \binom{k_1 + k_2 + k_3 - 3}{k_1 - 1} \binom{k_2 + k_3 - 2}{k_2 - 1} \\ &\times \widetilde{B}_{k_1+k_2+k_3+k_4-2}.\end{aligned}$$

(2) *If* $k_1 + k_2 + k_3 + k_4 \not\equiv 0 \pmod 2$, *then*

$$\frac{I_{k_1,k_2,k_3,k_4}(1)}{k_1!k_2!k_3!k_4!} = 0.$$

Remark 5.5. Our formula in Theorem 5.4 is equivalent to (5) in Carlitz [3, p. 361].

Letting $k_1 = 0$ and $k_2 = k, k_3 = l, k_4 = m$ in Theorem 5.4, we obtain the following formula for the integral of the product of three Bernoulli numbers:

Corollary 5.6. *For all* $k, l, m \geq 1$,

$$I_{k,l,m}(1) = \begin{cases} (-1)^{m+1} k!l!m! \sum_{a=0}^{k+l} \left[\binom{a}{l-1} + \binom{a}{k-1}\right] \widetilde{B}_{m+a+1} \widetilde{B}_{k+l-a-1} & \\ \qquad\qquad\qquad\qquad\qquad\qquad \textit{if } k+l+m \equiv 0 \pmod 2 \\ 0 \qquad\qquad\qquad\qquad\qquad\qquad\; \textit{if } k+l+m \not\equiv 0 \pmod 2. \end{cases}$$

REFERENCES

[1] T. Agoh and K. Dilcher, *Convolution identities and lacunary recurrences for Bernoulli numbers*, J. Number Theory **124** (2007) 105–122.

[2] T. Agoh and K. Dilcher, *Integrals of products of Bernoulli polynomials*, J. Math. Anal. Appl. **381** (2011) 10–16.

[3] L. Carlitz, *Note on the integral of the product of several Bernoulli polynomials*, J. London Math. Soc. **34** (1959) 361–363.

[4] K. Dilcher, *Sums of products of Bernoulli numbers*, J. Number Theory **60** (1996) 23–41.

[5] O. Espinosa and V.H. Moll, *The evaluation of Tornheim double sums. I*, J. Number Theory **116** (2006) 200–229.

[6] M. Mikolas, *Integral formulae of arithmetical characteristics relating to the zeta-function of Hurwitz*, Publ. Math. Debrecen **5** (1957) 44–53.

[7] L. J. Mordell, *Integral formulae of arithmetical character*, J. London Math. Soc. **33** (1958) 371–375.

[8] N. Nielsen, *Traite elementaire des nombres de Bernoulli*, Gauthier-Villars, Paris, 1923.

[9] N. E. Nörlund, *Vorlesungen uber Differenzenrechnung*, Springer-Verlag, Berlin, 1924.

[10] J. C. Wilson, *On Franel-Kluyver integrals of order three*, Acta Arith. **66** (1994) 71–87.

NATIONAL INSTITUTE FOR MATHEMATICAL SCIENCES, DORYONG-DONG, YUSEONG-GU, DAEJEON 305-340, SOUTH KOREA
E-mail address: `daeyeoul@nims.re.kr`

DEPARTMENT OF MATHEMATICAL SCIENCES, KOREA ADVANCED INSTITUTE OF SCIENCE AND TECHNOLOGY (KAIST), DAEJEON 305-701, SOUTH KOREA
E-mail address: `husu@kaist.ac.kr`

DIVISION OF CULTURAL EDUCATION, KYUNGNAM UNIVERSITY, CHANGWON 631-701, SOUTH KOREA
E-mail address: `mskim@kyungnam.ac.kr`